

High-Content Analysis of Chelidonine and Berberine from Iranian *Chelidonium majus* L. Ecotypes in Different Ontogenetical Stages Using Various Methods of Extraction

H. Pourmazaheri¹, B. Baghban Kohnerouz^{1*}, N. Khosravi Dehaghi², M. R. Naghavi³, E. Kalantar⁴, E. Mohammadkhani⁵, and M. Omidi³

ABSTRACT

Chelidonium majus is a perennial plant of the Papaveraceae family. This plant has been known as a rich source of isoquinoline alkaloids, chelidonine, and berberine, which are pharmaceutically important for their anti-cancerous activities. In the current study, four extraction techniques were compared in terms of their yield potential for chelidonine and berberine. Afterwards, High Performance Liquid Chromatography (HPLC) with a photodiode array type of UV/VIS detector was used for the detection of chelidonine and berberine from leaves and roots of five ecotypes of *C. majus* during various ontogenetical stages. Based on our results, ultrasonic procedures and refluxing were the best techniques for extraction of these alkaloids. HPLC results inferred that chelidonine and berberine content of ecotypes belonging to the Northern provinces of Iran, i.e. Mazandaran (IBRCP1006619) and Gorgan (IBRCP1006625), were higher than the other ecotypes. Generally, the roots of the *C. majus* were the most suitable organ for extraction of chelidonine at the generative stage, while at the vegetative stage, leaves are the most suitable organ for extraction of berberine.

Keywords: Extraction techniques, Generative stage, Isoquinoline alkaloids, Vegetative stage.

INTRODUCTION

Chelidonium majus L. is a perennial plant from the family of Papaveraceae. It is distributed in Europe and Western Asia and also an introduced species in Northern America. Commonly, this plant is known as celandine, greater celandine, celandine poppy, elon-wort, felon-wort, rock poppy, swallow-wort, and tetter-wort. The plant is highly toxic due to presence of various secondary

metabolites, but has been adequately used in traditional medicines (Monavari *et al.*, 2012).

Pharmacological properties of *C. majus* include anti-viral (Gerencer *et al.*, 2006), anti-bacterial (Miao *et al.*, 2011), anti-fungal (Hou *et al.*, 2013), anti-protozoal (Kim *et al.*, 2012), radioprotective (Song *et al.*, 2003), anti-inflammatory (Park *et al.*, 2011), anti-Alzheimer (Cahlikova *et al.*, 2010), anti-cancer (Moussa *et al.*, 2007), hepatoprotective (Biswas *et al.*, 2008), and natriuretic and antidiuretic (Koriem *et al.*, 2013). An array of

¹ Department of Plant Breeding and Biotechnology, College of Agriculture, University of Tabriz, Tabriz, Islamic Republic of Iran.

* Corresponding author; e-mail: baghban@tabrizu.ac.ir

² Department of Pharmacognosy, Faculty of Pharmacy, Alborz University of Medical Sciences, Karaj, Islamic Republic of Iran.

³ Department of Agronomy and Plant Breeding, College of Agriculture, University of Tehran, Karaj, Islamic Republic of Iran.

⁴ Department of Microbiology and Immunology, Faculty of Medicine, Alborz University of Medical Science, Karaj, Islamic Republic of Iran.

⁵ Iranian Biological Resource Center, Karaj, Islamic Republic of Iran.



secondary metabolites present in *C. majus* is responsible for its therapeutic properties. One of the most common secondary metabolites in *C. majus* is alkaloids. Various alkaloids metabolized by *C. majus* are chelidone, berberine, sanguinarine, coptisine, chelerythrine, protopine, etc. (Gilca *et al.*, 2010). In addition to alkaloids, other secondary metabolites identified from this plant include flavonoids, saponins, vitamins (e.g. vitamins A and C), mineral elements, sterols, acids and their derivatives (Kopytko *et al.*, 2005).

Two most pharmacologically important alkaloids identified from *C. majus* are chelidone and berberine. Chelidone is a benzophenanthridine alkaloid. This tertiary alkaloid ($C_{20}H_{19}NO_5$, molecular weight: 353.38 g mL⁻¹) has an anticancer effect (El-Readi *et al.*, 2013). On the other hand, berberine ($C_{20}H_{18}NO_4$, molecular weight: 336.37 g mL⁻¹) is an isoquinoline alkaloid having anti-cancer activity (Kemeny-Beke *et al.*, 2006; Mahata *et al.*, 2011). Several studies have been conducted on the quantification of chelidone and berberine from *C. majus* extracts (Kulpa *et al.*, 2011; Citoglu *et al.*, 2009; Gu *et al.*, 2010).

The concentration of alkaloids in plants is usually low, ranging between few percent (about 10 to 25%) and heterogeneously distributed within the plant tissue. Hence, adequate extraction methods are important for better recovery of alkaloids from a plant sources. Several studies have been published on the extraction of isoquinoline alkaloids, chelidone and berberine from *C. majus*. Among the published protocol four methods of extraction were applied for *C. majus* in order to compare extraction procedures and identify the best method which shows the highest rate of chelidone and berberine that save greater amount of anti-cancer agents in this plant. The salient features of these protocols are higher yield of alkaloids due to the used solvents that interact with the polar groups of alkaloids, faster extraction time, and lower operation costs.

The aim of this study was to compare four different optimized methods for extraction of

chelidone and berberine from *C. majus*, for selecting the most efficient extraction for recovering these alkaloids. In addition, we aimed to apply HPLC to quantify the chelidone and berberine content of five Iranian *C. majus* ecotypes, using different plant tissues at various ontogenetical stages. Another objective was to determine the richest source of these alkaloids among the selected ecotypes, to be used for future studies.

MATERIALS AND METHODS

Chemicals and Reagents

Standards for the alkaloids, chelidone (CAS No. 476-32-4; Lot No.BCBN1934V) and berberine (CAS No. 633-65-8; Lot No. SLBG1303V) were purchased from Sigma-Aldrich (St Louis, MO, USA). Methanol and acetonitrile (HPLC grade), n-Hexane, formic acid, hydrochloric acid, chloroform, ammonia (25%), citric acid, and ammonium formate in all experiments were purchased from Merck (Darmstadt, Germany). Deionized water was obtained and purified using a Millipore Milli-Q Plus water treatment system (Millipore Bedford Corp., Bedford, MA, USA).

Preparation of Standard Solutions

A stock solution containing 1 mg mL⁻¹ of each analyte was prepared in methanol (Merck, Germany) and stored at 4°C. The working standard solutions were made by dilution of the stock solution in methanol, and then filtered by 0.2- μ m membrane. Working solutions were stored at 4°C before injection into the HPLC.

Plant Material

Seeds of *C. majus* were collected from five different regions of the northern Iran (Table 1). The collections were submitted to the herbarium of the Iranian Biological Resources

Table 1. Regions where seeds of *C. majus* were collected.

Average temperature (°C)	Longitude coordinates (E)	Latitude coordinates (N)	Height (m)	Regions	Accession no
27	52 17' 0.9"	36 35' 15.1 "	1	Mazandaran; Mahmudabad- Amol	IBRCP1006619
22	52 57' 19.9"	35 52' 37.2"	1901	Mazandaran; Firoozkooh-Sari	IBRCP1006622
25	52 54' 32.9"	36 14' 26.9"	340	Mazandaran	IBRCP1006623
25	54 28' 52.9"	36 45' 23.4"	605	Gorgan	IBRCP1006625
29	54 7' 20.6"	36 46' 31.2"	34	Gorgan	IBRCP1006626

Center (IBRC), Alborz, Iran, under voucher number e.g., IBRCP1006619, IBRCP1006622, IBRCP1006623, IBRCP1006625, IBRCP1006626 that all ecotypes have similar morphology with the same age.

The seeds were cultivated in the greenhouse facilities of IBRC from November 2015 till March 2016. The temperature at the facility was maintained between 20-25°C and plants were irrigated regularly.

The aerial parts and the roots for the different ecotypes of *C. majus* were collected randomly at two different stages, i.e. vegetative and generative. The plant materials were cleaned with the deionized water and dried in the oven at 40°C, then, powdered (20 mesh) for further experiments.

Extraction Methods

In this study, based on the previous researches (Lee *et al.*, 2005; Artamonova and Kurkin, 2008; TalebiKouyakh *et al.*, 2008; Jahansooz *et al.*, 2008; Baghalian *et al.*, 2008; Baghalian *et al.*, 2010; Citoglu *et al.*, 2009; Sarkozi *et al.*, 2006; Gu *et al.*, 2010), the yield potential of four extraction methods for chelidonine and berberine were compared. These methods included: (1) Reflux with methanol (Ghanavi *et al.*, 2013); (2) Ultrasonic procedure with water-methanol-HCl (Zhou *et al.*, 2012); (3) Ultrasonic procedure with methanol-HCl

(Kursinszki *et al.*, 2006), and (4) Shaking incubator with citric acid (Borghini *et al.*, 2015).

High Performance Liquid Chromatography (HPLC) Analysis

HPLC analysis was carried out using Agilent 1200 series (Walbronn, Germany). HPLC system consisted of a G1312B binary pump, a G1376A capillary pump, G1330B FC/ALS, G1379B Degasser, and G1377Amicrowips. Chemstation software was used for data acquisition, processing and reporting. Compounds were separated on a 5 µm ZORBAX Eclipse XDBC18 reversed-phase column (150 mm×4.6 mm id; Agilent Technology, Germany). The column temperature was 30°C and the injection volume was 20 µL (in triplicate).

In order to simultaneously determine chelidonine and berberine, a gradient program with two solvent systems including A, water (0.1% formic acid), and B, acetonitrile, was applied. Separation was performed with 20% B at the beginning, which gradually changed to 90% B within 40 minutes, remaining 5 minutes at this point for re-equilibration. The flow rate of the mobile phase was 0.5 mL min⁻¹. The photodiode detector was scanned from 210 to 400 at 2-nm intervals. The chromatographic profile was recorded at 280 nm. Retention times and UV-visible



absorption spectra of the samples were compared with the standards.

Statistical Analysis

All tests were performed in triplicate using factorial experimental design. Data were expressed as mean±SD values, while significance (P values less than 0.05 to 0.01) of the differences between the mean values was determined by Duncan's multiple range test using Statistical Analysis System Ver. 9.1 (SAS) software.

RESULTS AND DISCUSSION

Calibration Curves

Calibration curves for chelidonium were drawn at 2,500, 5,000, 10,000, 25,000 ng mL⁻¹, whereas in case of berberine it was 60, 125, 250, 500, 1,000, 2,500, 5,000 ng mL⁻¹. Linear regression and linear range of each alkaloid are shown in Table 2. All calibration curves showed the best-fitting linear regression ($r \geq 0.999$) within tested ranges. Twenty µL of each standard solution in triplicate was injected on to the HPLC column and absorbance was recorded at 280 nm. The concentrations of chelidonium and berberine in samples were calculated from their peak areas by use of the calibration curve.

HPLC Analysis

HPLC is commonly used for separation of metabolites from *C. majus* extract using both normal and Reverse Phase (RP) columns

(Suchomelova et al., 2007; Taborska et al., 1994; Bugatti et al., 1987; Zuo et al., 2008).

Application of HPLC for separation of chelidonium and berberine requires some modifications to obtain fast and efficient protocol for high recovery within a short run time (less than 20 min). In the present study, due to the effect of polar functional groups and chromatographic behavior of chelidonium and berberine induced by non-polar HPLCC18 column; acetonitrile-formic acid was selected as the most convenient elution mobile phase following Ghanavi et al. (2015).

Typical Chromatogram for chelidonium and berberine mixed standards (5,000 ng mL⁻¹) are shown in Figure 1. For each sample, analysis was carried out in triplicate. Chromatograms obtained for chelidonium and berberine extracts from leaves and roots of five ecotypes of *C. majus* during vegetative and generative stages are shown in Figures 2-5. As depicted, two alkaloids separated within 12 minutes and data represent the means±standard deviation of three independent experiments (Table 4). The ANOVA of the different tissues and growth stages data of *C. majus* in five ecotypes is shown in Table 5.

Extraction Methods

Selection of extraction procedures and solvents depends on several factors, which includes the physicochemical nature of secondary metabolite, nature of plant material (fresh parts, dried parts) and their particle size (Yadav et al., 2003). As the

Table 2. Linear regression data of chelidonium and berberine.^a

Linear range (ng mL ⁻¹)	r	Regression equations	Alkaloid
2500-25000	0.999	$Y = 35.80x - 17.22$	Chelidonium
60-5000	0.999	$Y = 0.125x + 1.53$	Berberine

^a $Y = Ax + B$, Y is peak area; x is concentration of the alkaloids (ng mL⁻¹), r is the correlation coefficient of the equation.




Figure 1. Chromatogram of Chelidone (CHE) and Berberine (BER) mixed standards.




Figure 2. Separated chelidone and berberine chromatograms from roots of five Iranian *C. majus* ecotypes at vegetative stage.




Figure 3. Separated chelidone and berberine chromatograms from roots of five Iranian *C. majus* ecotypes at generative stage.




Figure 4. Separated chelidone and berberine chromatograms from leaves of five Iranian *C. majus* ecotypes at vegetative stage.

chemical property of a solvent is directly correlated to the final yield, solvents such as methanol and its mixtures (methanol-water, acidic methanol with HCl) along with acetic

acid and other solvents were selected on the basis of polarity of alkaloids (Madziga *et al.*, 2010; Yalavarthi and Thiruvengadarajan, 2013).




Figure 5. Separated chelidone and berberine chromatograms from leaves of five Iranian *C. majus* ecotypes at generative stage.

Our results showed that ultrasonic and reflux methods were the most efficient techniques for extracting chelidone and berberine, respectively. Concentration of chelidone and berberine obtained using different extraction methods are presented in Table 3.

Among the five Iranian *C. majus* ecotypes investigated in this study, the final yield of chelidone and berberine substantially varied between different parts of the plant and growth stages (Table 4). Among ecotypes, we recovered the highest amount of chelidone from IBRCP1006619 (Mahmudabad-Amol, Mazandaran), followed by IBRCP1006622 (Firoozkooh-Sari, Mazandaran) and IBRCP1006623 (Mazandaran). Ecotypes from Gorgan, *i.e.*

IBRCP1006625 and IBRCP1006626 showed a great variation in concentration of chelidone in terms of different plant tissues and different ontogenetical stages. Irrespective of the ecotypes, chelidone concentrations in the roots increased with the transition of the plant from vegetative stage to generative stage. While this relationship was reverse in the case of the leaf tissues. For all ecotypes, chelidone concentration in the roots were higher than that of the leaves. In terms of plant growth stages, the chelidone concentration was relatively higher in the generative stage than in the vegetative and this is applicable to all ecotypes.

In terms of berberine content, ecotypes from Gorgan had significantly more of this alkaloid than Mazandaran. IBRCP1006625 (Gorgan) in vegetative stage produced high levels of berberine, followed by IBRCP1006623, IBRCP1006622 and IBRCP1006619 (Mazandaran), and lastly IBRCP1006626 (Table 4). The concentration of berberine was higher in the leaves than in the roots for all ecotypes. In term of growth stages, the vegetative stage had a relatively higher concentration of berberine than in the generative stage (Table 4). Berberine concentrations in all ecotypes decreased during the transition of the plant from vegetative to the generative stage.

By statistically comparing our data, it revealed that the ecotypes with the higher amount of berberine in leaves showed lower levels of chelidone in the roots. Among the ecotypes from Mazandaran, IBRCP1006619 produced high levels of chelidone (19.13 ± 0.15 mg g⁻¹ of dry weight) at the generative stage in root and the ecotype of Gorgan, IBRCP1006625

Table 3. Concentration of isolated chelidone and berberine with different extraction methods.

Extraction method	Alkaloids content	
	Chelidone (mg g ⁻¹ DW)	Berberine (μ g mg ⁻¹ DW)
Refluxing with methanol	6.57 ± 0.11	0.90 ± 0.09
Ultrasonic with water-methanol-HCl	10.99 ± 0.14	0.42 ± 0.02
Ultrasonic with methanol-HCl	12.04 ± 0.01	0.43 ± 0.03
Shaking incubator with Acidic condition on dried parts	6.85 ± 0.11	0.05 ± 0.02

Table 4. Isolated chelidonine (mg g^{-1} DW) and berberine ($\mu\text{g mg}^{-1}$ DW) contents from leaves and roots of five ecotypes of *C. majus* during vegetative and generative stages.

Growth stages		Metabolite	Ecotype	Organ
Generative stage	Vegetative stage			
19.13± 0.15	15.52± 0.28	CHE	IBRC P1006619	Root
0.29± 0.01	0.53± 0.02	BER		
14.45± 0.28	12.87± 0.31	CHE	IBRC P1006622	
0.86±0.02	1.65±0.01	BER		
13.26± 0.15	12.5 ±0.30	CHE	IBRC P1006623	
0.28± 0.02	0.42 ± 0.01	BER		
12.21± 0.09	11.93± 0.24	CHE	IBRC P1006625	
0.10± 0.02	0.21± 0.01	BER		
10.93± 0.15	9.15±0.05	CHE	IBRC P1006626	
0.37± 0.01	0.54± 0.08	BER		
6.52± 0.24	5.08± 0.02	CHE	IBRC P1006619	Leaf
0.43± 0.02	1.65± 0.03	BER		
11.99± 0.23	6.33± 0.21	CHE	IBRC P1006622	
0.45± 0.01	2.00±0.02	BER		
14.45± 0.22	8.17± 0.04	CHE	IBRC P1006623	
0.51± 0.01	2.83±0.01	BER		
15.23± 0.14	9.60± 0.05	CHE	IBRC P1006625	
0.56± 0.01	3.98 ± 0.02	BER		
15.58± 0.11	10.96± 0.14	CHE	IBRC P1006626	
0.42± 0.01	1.53±0.01	BER		

Table 5. Analysis Of Variance (ANOVA) of the different tissues and growth stages data of *C. majus* in five ecotypes.

Source of variation	Degrees of freedom	MS
Tissue	1	117.91
Ecotype	4	2.88
Growth stage	1	150.07
Tissue×Ecotype	4	93.75
Tissue×Growth stage	1	36.49
Ecotype×Growth stage	4	0.47
Tissue×Ecotype×Growth stage	4	7.56
Error	38	0.03
Coefficient of variation	1.52	

produced highest levels of berberine ($3.98 \pm 0.02 \mu\text{g mg}^{-1}$ of dry weight) at vegetative stage in leaf. Hence, it is quite evident that the metabolisms for these alkaloids change between the developmental stages of the plant and between tissues.

In previous study, Ghanavi *et al.* (2013) quantified the isoquinoline alkaloids content of *C. majus* using refluxing technique. Their study showed berberine content of *C. majus* stem was $1.28 \mu\text{g mg}^{-1}$. Sarkozi *et al.* (2006)

investigated the chelidonine content from the leaf and root in *C. majus*, and their result is consistent with our study. They observed a significant difference in chelidonine content in various organs of *C. majus*. The chelidonine concentration in roots and leaves was more than in other organs. They recovered 0.18 and 3.76 mg of chelidonine from dry weight of leaf and root, respectively. Gu *et al.* (2010) investigated the chelidonine content from the roots of *C.*



majus collected from different regions in China and Mongolia by Ultra-Performance LC (UPLC) method with photodiode array detection. The maximum chelidone content from these two regions were about 2.34 ± 0.00 and 2.87 ± 0.05 mg, respectively. They recovered a low berberine concentration from all samples (averaged 0.13 mg). In another study, the differences of chelidone and berberine contents in *C. majus* extracts from 14 different geographic areas in China and showed that the ranges of chelidone and berberine recovered were 514 to 1,012 mg kg⁻¹ and 499 to 1,161 mg kg⁻¹, respectively (Zhou *et al.*, 2012).

In our study, we reconfirmed ultrasonic and reflux as better methods than shaking incubator for extraction of chelidone and berberine from *C. majus*. The advantages of ultrasonication compared to the reflux and shaking incubator are requirement of shorter extraction time and temperature, conserving the chemical structure of the metabolites. In contrast, shaking needs longer extraction time and higher temperature giving less yield due to decomposition of the metabolites (Arulpriya *et al.*, 2013). However, reflux is another efficient method that in some cases was superior than ultrasonic extraction (Lee *et al.*, 2013), but simplicity of the apparatus, faster operational time, and ability to tandem process multiple samples (Porevsky *et al.*, 2014; Manika *et al.*, 2013) makes ultrasonication the preferred extraction method. For the first time, we recovered the highest amount of chelidone and berberine using ultrasonic and reflux techniques coupled with developmental stages of the *C. majus*. We believe that the higher yield of the alkaloids in our study was outcome of the different ontogenetical stages of the plant we investigated.

CONCLUSIONS

In the present study, chelidone and berberine contents from five ecotypes of *C. majus* were assessed from roots and leaves

at different developmental stages using four different extraction methods. Since even slight increase in the amount of chelidone and berberine is more important than optimizing a new extraction method; in this study, the yield potential of four already published methods were compared to reach the highest amount of these two alkaloids. Based on our results, the root is the most suitable organ for extraction of chelidone at the generative stage. In the case of berberine, leaves at the generative stage are the most suitable organ for extraction. Generally, the ecotypes from Mazandaran (IBRCP1006619) and Gorgan (IBRCP1006625) showed a higher concentration of chelidone and berberine, respectively. According to the results, we determined the richest source of these alkaloids among the selected ecotypes, which can be used for RNA-sequencing to discover a potential connection between gene expression and content variation for chelidone and berberine, using differential regulatory mechanisms underlying secondary metabolic pathways in two different tissues of *C. majus*.

ACKNOWLEDGEMENTS

The authors would like to acknowledge University of Tabriz and Alborz University of Medical Sciences for financial support of this research.

REFERENCES

1. Artamonova, E. S. and Kurkin, V. A. 2008. Developing Methods for Qualitative and Quantitative Analysis of *Chelidonium majus* Herbs. *Pharm. Chem. J.*, **42(11)**: 633-636.
2. Arulpriya, P. and Lalitha, P. 2013. Evaluation of Different Extraction Methods for Optimization of Extraction of Aerial Roots of *Rhaphidophora aurea* Entwined over Two Diverse Host Trees. *Int. J. Chem. Tech. Res.*, **5(5)**: 2173-2176.
3. Baghalian, K. Haghiry, A. Naghavi, M. R. and Mohammadi A. 2008. Effect of Saline

- Irrigation Water on Agronomical and Phytochemical Characters of Chamomile (*Matricaria recutita* L.). *Scientia Hort.*, **116**:437-441.
4. Baghalian, K., Maghsodi, M. and Naghavi, M. R. 2010. Genetic Diversity of Iranian Madder (*Rubia tinctorum*) Populations Based on Agro-Morphological Traits, Phytochemical Content and RAPD Markers. *Ind. Crop. Prod.*, **31**: 557-562.
 5. Biswas, S. J., Bhattacharjee, N. and Khudabukhsh, A. R. 2008. Efficacy of a Plant Extract (*Chelidonium majus* L.) in Combating Induced Hepatocarcinogenesis in Mice. *Food Chem. Toxicol.*, **46** (5): 1474-1487.
 6. Borghini, A. Pietra, D., Trapani, C., Madau, P., Lubinu, G. and Bianucci, A. 2015. Data Mining as a Predictive Model for *Chelidonium majus* Extracts Production. *Ind. Crop. Prod.*, **64**: 25-32.
 7. Bugatti, C., Colombo, M.L. and Tome, F. 1987. High-Performance Liquid Chromatographic Separation of Quaternary Alkaloids of *Chelidonium majus* L. Root. *J. Chromatogr.*, **393**: 312-316.
 8. Cahlikova, L., Opletal, L., Kurfurst, M., Macakova, K., Kulhankova, A. and Hostalkova, A. 2010. Acetylcholinesterase and Butyrylcholinesterase Inhibitory Compounds from *Chelidonium majus* (Papaveraceae). *Nat. Prod. Commun.*, **5** (11): 1751-1754.
 9. Ciric, A., Vinterhalter, B., Savikin-Fodulovic, K., Sokovic, M. and Vinterhalter, D. 2008. Chemical Analysis and Antimicrobial Activity of Methanol Extracts of Celandine (*Chelidonium majus* L.) Plants Growing in Nature and Cultured *In Vitro*. *Arch. Biol. Sci.*, **60**(1): 7-8.
 10. Citoglu, G., Ozbek, H., Acikara, O. and Gacs, E. 2009. Isolation of Chelidonine as an Analgesic Compound from *Chelidoniummajus* L. *J. Fac. Pharm. (Ankara)*, **38**(1): 9-16.
 11. El-Readi, M. Z. Eid, S. Ashour, M. L. Tahrani, A. and Michael, W. 2013. Modulation of Multidrug Resistance in Cancer Cells by Chelidonine and *Chelidonium majus* Alkaloids. *Phytomedicine*, **20**: 282-294.
 12. Gerencer, M. Turecek, P. L. Kistner, O. Mitterer, A. Savidis-Dacho, H. and Barrett, N. P. 2006. *In Vitro* and *In Vivo* Anti-Retroviral Activity of the Substance Purified from the Aqueous Extract of *Chelidonium majus* L. *Antiviral. Res.*, **72**(2): 153-156.
 13. Ghanavi, Z. Eslami, Z. Mollayi, S. NaghdiBadi, H. and Babaei, A. 2013. Quantification of Isoquinoline Alkaloids Content in Stem of Celandine (*Chelidonium majus*) from North of Iran. *Intl. J. Agron. Plant. Prod.*, **4**(8): 2039-2045.
 14. Ghanavi, Z., Abdousi, V., Samadian-Sarbangholi, V., Mollayi, S., Babaei, A. and Ghassempour, A. 2015. Effect of Environmental Factors on Sanguinarine and Berberine Levels in Root of *Chelidonium majus* by HPLC- PDA/MS Method. *J. Chem. Pharm. Res.*, **7**(6): 15-21.
 15. Gilca, M., Gaman, L., Panait, E., Stoian, I. and Atanasiu, V. 2010. *Chelidonium majus*-an Integrative Review: Traditional Knowledge versus Modern Findings. *Forsch Komplementmed*, **17**(5): 241-248.
 16. Gu, Y., Qian, D., Duan, J., Wang, Z., Guo, J., Tang, Y. and Guo, Sh. 2010. Simultaneous Determination of Seven Main Alkaloids of *Chelidonium majus* L. by Ultra-Performance LC with Photodiode-Array Detection. *J. Sep. Sci.*, **33**(8): 1004-1009.
 17. Hou, Z., Yang, R., Zhang, C., Zhu, L. F., Miao, F., Yang, X. J. and Zhou, L. 2013. 2-(Substituted Phenyl)-3,4-Dihydroisoquinolin-2-Iums as Novel Antifungal Lead Compounds: Biological Evaluation and Structure-Activity Relationships. *Mol.*, **18** (9): 10413-10424.
 18. Jahansooz, F. Ebrahimzadeh, H. Najafi, A. A. Naghavi, M. R. Kouyakh, E. T. and Farzaneh, H. 2008. Composition and Antifungal Activity of the Oil of *Ferula gummosa* Samples from Iran. *J. Essent. Oil. Bear. Pl.*, **11**: 284-291.
 19. Kemeny-Beke, A., Aradi, J., Damjanovich, J., Beck, Z., Facsko, A., Berta, A. and Bodnar, A. 2006. Apoptotic Response of Uveal Melanoma Cells upon Treatment with Chelidonine, Sanguinarine and Chelerythrine. *Cancer Lett.*, **237**: 67-75.
 20. Kim, D. S., Kim, S. J., Kim, M. C., Jeon, Y. D., Um, J. Y. and Hong, S. H. 2012. The Therapeutic Effect of Chelidonic Acid on Ulcerative Colitis. *Biol. Pharm. Bull.*, **35**(5): 666-671.
 21. Kopytko, Y. F., Dargaeva, T. D., Sokolskaya, T. A., Grodnitskaya, E. I. and Kopnin, A. A. 2005. New Methods for the Quality Control of a Homeopathic Matrix Tincture of Greater



- Celandine. *Pharm. Chem. J.*, **39(11)**: 603-609.
22. Koriem, K. M., Arbid, M. S. and Asaad, G. F. 2013. *Chelidonium majus* Leaves Methanol Extract and Its Chelidonine Alkaloid Ingredient Reduce Cadmium-Induced Nephrotoxicity in Rats. *J. Nat. Med.*, **67(1)**: 159-167.
 23. Kulpa, M., Braginab, O., Kogermanb, P. and Kaljuranda, M. 2011. Capillary Electrophoresis with Led-Induced Native Fluorescence Detection for Determination of Isoquinoline Alkaloids and Their Cytotoxicity in Extracts of *Chelidonium majus* L. *J. Chromatogr. A*, **1218(31)**: 5298-5304.
 24. Kursinszki, L., Sarkozi, A., Kery, A. and Szoke, E. 2006. Improved RP-HPLC Method for Analysis of Isoquinoline Alkaloids in Extracts of *Chelidonium majus*. *Chromatographia*, **63**: S131-S135.
 25. Lee, J., Shon, M. Y., Jang, D. S., Ha, T. J., Hwang, S. W., Nam, S. H. and Yang, M. S. 2005. Cytotoxic Isoquinoline Alkaloids from *Chelidonium majus* var. *Asiaticum*. *J. Appl. Biol. Chem.*, **48(4)**: 198-201.
 26. Lee, L. S., Lee, N., Kim, Y. H., Lee, C. H., Hong, S. P., Jeon, Y. W. and Kim, Y. E. 2013. Optimization of Ultrasonic Extraction of Phenolic Antioxidants from Green Tea Using Response Surface Methodology. *Mol.*, **18(11)**: 13530-13545.
 27. Madziga, H. A., Sanni, S. and Sandabe, U. K. 2010. Phytochemical and Elemental Analysis of *Acalypha wilkesiana* Leaf. *J. Am. Sci.*, **6(11)**: 510-514.
 28. Mahata, S., Bharti, A. C., Shukla, S., Tyagi, A., Husain, S. A. and Das, B. C. 2011. Berberine Modulates AP-1 Activity to Suppress HPV Transcription and Downstream Signaling to Induce Growth Arrest and Apoptosis in Cervical Cancer Cells. *Mol. Cancer*, **10(39)**: 1-14.
 29. Miao, F., Yang, X. J., Zhou, L., Hu, H. J., Zheng, F., Sun, X. D., Ding, D. M., Zhou, C. D. and Sun, W. 2011. Structural Modification of Sanguinarine and Chelerythrine and Their Antibacterial Activity. *Nat. Prod. Res.*, **25(9)**: 863- 875.
 30. Manika, N., Gupta, V. K., Verma, R. K., Darokar, M. P., Pandey, N. and Bagchi, G. D. 2013. Extraction Efficacy, Antibacterial Potential And Validation Of RP-HPLC Coupled With Diode Array Detection In *Holarrhena pubescens*. *Int. J. Res. Pharma. Sci.*, **4(8)**: 3020.
 31. Monavari, S. H., Shahrabadi, M. S., Keyvani, H. and Bokharaei-Salim, F. 2012. Evaluation of *In Vitro* Antiviral Activity of *Chelidonium majus* L. against Herpes Simplex Virus Type-1. *Afr. J. Microbiol. Res.*, **6(20)**: 4360- 4364.
 32. Moussa, S. Z., El-Meadawy, S. A., Ahmed, H. A. and Refat, M. 2007. Efficacy of *chelidonium majus* and Propolis against Cytotoxicity Induced by Chlorhexidine in Rats. *J. Biochem. Mol. Biol.*, **25**: 42-68.
 33. Park, J. E., Cuong, T. D., Hung, T. M., Lee, I., MinKyun, Na., Kim, J. C. Ryoo, S. W., Lee, J. H., Choi, J. S., Woo, M. H. and Min, B. S. 2011. Alkaloids from *Chelidonium majus* and Their Inhibitory Effects on LPS-Induced NO Production in RAW264.7 Cells. *Bioorg. Med. Chem. Lett.*, **21(23)**: 6960-6963.
 34. Porevsky, P. A., Ruiz, H. G. and Garciadiego, L. H. 2014. Comparison of Soxhlet Extraction, Ultrasonic Bath and Focused Microwave Extraction Techniques for the Simultaneous Extraction of PAH's and Pesticides from Sediment Samples. *Sci. Chromatogra*, **6(2)**:124-138.
 35. Sarkozi, A., Janicsak, G., Kursinszki, L. and Kery, A. 2006. Alkaloid Composition of *Chelidonium majus* L. Studied by Different Chromatographic Techniques. *Chromatographia*, **63**: S81-S86.
 36. Song, J. Y., Yang, H. O., Shim, J. Y., Han, Y. S., Jung, I. S. and Yun, Y. S. 2003. Radiation Protective Effect of an Extract from *Chelidonium majus*. *Int. J. Hematol.*, **78(3)**: 226-232.
 37. Suchomelova, J., Bochorakova, H., Paulova, H., Musil, P. and Taborska, E. 2007. HPLC Quantification of Seven Quaternary Benzo[c]phenanthridine Alkaloids in Six Species of the Family *Papaveraceae*. *J. Pharm. Biomed. Anal.*, **44**: 283-287.
 38. Taborska, E., Bochorakova, H., Paulova, H. and Dostal, J. 1994. Separation of Alkaloids in *chelidonium majus* by Reversed-Phase HPLC. *Planta. Med.*, **60(4)**: 380-381.
 39. TalebiKouyakhhi, E., Naghavi, M. R. and Alayhs, M. 2008. Study of the Essential Oil Variation of *Ferula gummosa* Samples from Iran. *Chem. Nat. Compd.*, **44**: 124-126.
 40. Yadav, J. S., Reddy, B. V. S. and Premalatha, K. 2003. 1-Butyl-3-Methylimidazoliumtetrafluoroborate ([Bmim] BF) Ionic Liquid: A Novel and Recyclable

- Reaction Medium for the Synthesis of Vic-Diamines. *Adv. Synth. Catal.*, **345**: 948-952.
41. Yalavarthi, Ch. and Thiruvengadarajan, V. S. 2013. A Review on Identification Strategy of Phytoconstituents Present in Herbal Plants. *Int. J. Res. Pharma. Sci.*, **4(2)**: 123-140.
42. Zhou, Q., Liu, Y., Wang, X. and Di, X. 2012. Microwave-Assisted Extraction in Combination with Capillary Electrophoresis for Rapid Determination of Isoquinoline Alkaloids in *Chelidonium majus* L. *Talanta*, **99**: 932-938.
43. Zuo, J. L., Bai, L., Song, X., Gu, Y. and Zhao, C. 2008. Simultaneous Determination of Sanguinarine, Berberine and Chelerythrine in *Chelidonium majus* by RP-HPLC. *J. Pharm. Anal.*, **28**: 903-905.

بررسی مقادیر بالای کلیدونین و بربرین در اکوتیپ های ایرانی مامیران کبیر طی مراحل رشدی مختلف با استفاده از روش های مختلف استخراج

ه. پورمظاهری، ب. باغبان کهنه روز، ن. خسروی دهقی، م. ر. نقوی، ع. کلانتر،
 ا. محمدخانی، و م. امیدی

چکیده

مامیران کبیر (*Chelidonium majus*) گیاهی چندساله از خانواده خشخاش می باشد. این گیاه به علت دارا بودن منبع غنی از آلکالوئیدهای ایزوکوینولین، کلیدونین و بربرین شناخته شده است. اهمیت کلیدونین و بربرین از لحاظ دارویی به دلیل فعالیت ضدسرطانی آنهاست. در این مطالعه چهار روش استخراجی جهت دستیابی به بالاترین عملکرد آلکالوئیدهای کلیدونین و بربرین مورد مقایسه قرار گرفت. سپس کروماتوگرافی مایع با کارایی بالا دارای دکتور UV/VIS از نوع آرایه دیود حساس نسبت به نور جهت تشخیص کلیدونین و بربرین در بافت های برگ و ریشه پنج اکوتیپ مامیران در مراحل رشدی رویشی و زایشی به کار گرفته شد. طبق نتایج بدست آمده روش های استخراجی مبتنی بر اولتراسونیک و رفلکس، بهترین روش ها برای استخراج آلکالوئیدهای مذکور بود. نتایج کروماتوگرافی مایع با کارایی بالا نشان داد محتوای کلیدونین و بربرین به ترتیب در اکوتیپ های استان های شمالی ایران، مازندران (IBRCP1006619) و گرگان (IBRCP1006625) از سایر اکوتیپ ها بالاتر بود. در کل، ریشه بهترین اندام مناسب برای استحصال کلیدونین در فاز زایشی بود درحالی که برگ ها در فاز رویشی بهترین اندام مناسب جهت استحصال بربرین می باشد.