

Evaluating Allelopathic Effects of Some Plant Species in Tissue Culture Media as an Accurate Method for Selection of Tolerant Plant and Screening of Bioherbicides

E. Aryakia^{1,4*}, M. R. Naghavi², Z. Farahmand¹, and S. A. H. Shahzadeh Fazeli^{1,3}

ABSTRACT

Plant tissue culture technique could provide sterile and controllable condition in order to assay direct effect of different compounds on plant growth accurately. In this study, the effects of aqueous extracts prepared from roots and shoots of goosefoot (*Chenopodium album* L.), redroot pigweed (*Amaranthus retroflexus* L.), fennel (*Foeniculum vulgare*), and wormwood (*Artemisia absinthium* L.) were evaluated on the seed germination and growth criteria in tissue culture media. The fennel root extract, nearly without phenolic content and with low antioxidant activity, showed the most drastic allelopathic effect on goosefoot, especially at 100 mg mL⁻¹ concentration, which might be due to the presence of some substance potentially useful for biological control of goosefoot, an invasive weed. Goosefoot was resistant to extract of fennel shoot, wormwood root, and shoot, while fennel and radish (*Raphanus sativus* L.), at high concentration (100 mg mL⁻¹), were not resistant to the root and shoot extracts of both goosefoot and redroot pigweed. In response to allelopathic components, shoot:root ratio was increased, and more peroxidase and superoxide dismutase activity were detected in roots. There was no direct relationship between allelopathic effects with total phenolic content and antioxidant activity. In conclusion, our results showed that allelopathic effects of extracts on growth and biochemical criteria depended on both the concentration levels and the plant parts from which the aqueous extract was derived. Therefore, tissue culture media is an accurate and suitable tool to screen plants resistant to allelopathic components of weeds, and to identify high allelopathic plants as potential bioherbicide and invasive plant controller.

Keywords: Aqueous extract, Growth criteria, Invasive plant, Total phenolic content, Antioxidant activity.

INTRODUCTION

The phenomenon of allelopathy encompasses all types of direct and indirect chemical interactions among plants and microorganisms (Einhellig, 1995). Almost all kinds of plant species could produce allelochemicals that vary among plant species, plant part, and growing stages and

could be harmful and poisonous for one or more species (Sodaeizadeh *et al.*, 2009; Oueslati, 2003; Cheel *et al.*, 2012). Evaluating allelopathic effect of plant species could be important in different environmental conditions such as producing agricultural crops (Xuan *et al.*, 2012; Kato-Noguchi *et al.*, 2012), bioherbicides in weeds controlling strategies (Teerarak *et al.*,

¹ Plant Bank, Iranian Biological Resource Center (IBRC), ACECR, Tehran, Islamic Republic of Iran.

*Corresponding author; e-mail: aryakia@ibrc.ir or elyasaryakia@yahoo.com

² Department of Agronomy and Plant Breeding, University College of Agriculture and Natural Resources, University of Tehran, Karaj, Islamic Republic of Iran.

³ Department of Genetics, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Islamic Republic of Iran.

⁴ Department of Horticultural Sciences, Faculty of Agricultural Sciences, Vali-e-asr University of Rafsanjan, Islamic Republic of Iran.



2012; Tigrea *et al.*, 2012) and the effect of allelochemicals on biodiversity and establishing of invasive species (Valera-Burgos *et al.*, 2012; Inderjit, 1996; Tognetti and Chaneton, 2012). Research has shown that many factors including nutrition, humidity, radiation, temperature, pest and disease influence the allelopathy process (Einhellig, 1996). For example, microorganisms positively or negatively interfere with plant-plant allelopathy (Mishra and Nautiyal, 2011; Anaya *et al.*, 2013; Cipollini *et al.*, 2012) or could directly have negative (Goodal *et al.*, 2012; Elliott and Cheng, 1987) or positive (Mallik and Williams, 2008; Sturz and Christie, 2003) allelopathic effects on plant (Barazani and Friedman, 2001).

It seems that by control stressful environmental conditions, real effect of plant would be emerged. Plant tissue culture media can provide sterile and controllable conditions in which accurate and direct effect of different components such as organic materials (Aryakia and Hamidoghli, 2010; Siril and Joseph, 2013) or mineral elements (Bouman and Tiekstra, 2001) on growth could be assayed. The aims of this study were: 1) to evaluate allelopathic effects of some plant species on germination and growth criteria using tissue culture media, as a new method for identifying plant with high allelopathic effect (probably as bioherbicides and invasive plants controllers); and 2) to identify resistant plants to allelopathic components.

MATERIALS AND METHODS

Plant Material and Preparation of Aqueous Extracts

Seeds, shoots, and roots of goosefoot (*Chenopodium album* L.), redroot pigweed (*Amaranthus retroflexus* L.), fennel (*Foeniculum vulgare*), wormwood (*Artemisia absinthium* L.) and radish (*Raphanus sativus* L.) were separately collected at reproductive stage in the

glasshouse of Iranian Biological Resource Center, Alborz province, Iran. The surfaces of roots were washed with distilled water and were desiccated between the sheets of paper for one minute. Then, 250 g fresh matter of each part was separately cut into one-centimeter pieces, and shaken in 1000 mL double distilled water at 125 rpm at a temperature of 25 °C for 48 hours. Extracts were separated from the residue using two layer Whatman filter paper No. 42 (125 mm) and centrifuged at 5,000 rpm for 30 minutes. Supernatant was used as 100% stock solution (250 mg mL⁻¹) and stored at -20 °C.

Preparation of Tissue Culture Medium

To evaluate allelopathic effects of plant extracts on tissue culture media, modified MS medium (Murashige and Skoog, 1962) containing macronutrients: ammonium nitrate (NH₄NO₃) 1.650 mg L⁻¹, calcium chloride (CaCl₂·2H₂O) 440 mg L⁻¹, magnesium sulfate (MgSO₄·7H₂O) 370 mg L⁻¹, potassium phosphate (KH₂PO₄) 170 mg L⁻¹, potassium nitrate (KNO₃) 1.900 mg L⁻¹, and micronutrients: boric acid (H₃BO₃) 6.2 mg L⁻¹, cobalt chloride (CoCl₂·6H₂O) 0.025 mg L⁻¹, cupric sulfate (CuSO₄·5H₂O) 0.025 mg L⁻¹, ferrous sulfate (FeSO₄·7H₂O) 27.8 mg L⁻¹, manganese sulfate (MnSO₄·4H₂O) 22.3 mg L⁻¹, potassium iodide (KI) 0.83 mg L⁻¹, sodium molybdate (Na₂MoO₄·2H₂O) 0.25 mg L⁻¹, zinc sulfate (ZnSO₄·7H₂O) 8.6 mg L⁻¹, Na₂EDTA·2H₂O 37.2 mg L⁻¹ were used. This medium was supplemented with vitamins including inositol 100 mg L⁻¹, niacin 0.5 mg L⁻¹, pyridoxine-HCl 0.5 mg L⁻¹, thiamine-HCl 0.1 mg L⁻¹ and sucrose 30 g L⁻¹. The pH of the medium was adjusted to 5.8 before heat sterilization (at 121 °C for 15 minutes) by 0.1N KOH and then 8 gr L⁻¹ agar was added. With the aim of conserving heat-sensitive allelochemicals, extracts were added after autoclave. The pH of the extracts was adjusted to 5.8 and was sterilized by 0.2 micron-rated filter membrane. Later, 30 mL of the media was added to each 300 mL flask.

Disinfection and Seed Culture

Small and damaged seeds were removed before starting disinfection. Seeds were washed with tap water, rinsed with 70% (v/v) ethanol for 30 seconds and then were disinfected by sodium hypochlorite (NaClO) 1.5% (w/v) for 20 minutes. Seeds were washed with sterile distilled water several times under laminar flow and cultured in the media. The culture condition (temperature $24\pm 2^\circ\text{C}$, 16 hours photoperiod using cool white fluorescent light, and 3,000–4,000 lux light intensity) were maintained throughout the study. Germination, growth, and biochemical criteria were measured during 60 days of culture initiation.

Methanolic Extract Preparation

To determine total phenolic content and antioxidant activity, plants root and shoot were collected from the fields at the reproductive stage and dried at 40°C for 48 hours. Five hundred mg of dry weight for each sample was powdered and shaken for 24 hours in 10 mL of 80% (v/v) methanol. Extracts were then separated from the residue using Whatman No.1 filter paper and then centrifuged at 4,000 rpm for 20 minutes. The supernatant was used to determine the total phenolic content and antioxidant activity.

Determination of Total Phenolic Content

The amount of total phenolic was determined according to the method of Singleton and Orthofer (1999), which used Folin-Ciocalteu reagent and gallic acid as standard. Thirty μL of methanolic extract of root or shoot was diluted to the volume of 500 μL by distilled water, and then 500 μL of undiluted Folin-Ciocalteu reagent were added and mixed. The mixture was allowed to stand for a 1 min period, followed by the addition of 500 μL of 20% (w/v) sodium

carbonate (Na_2CO_3). After standing in the dark and room temperature for 120 minutes, the absorbance was read at 730 nm using a UV-Vis spectrophotometer. The total phenolic content was expressed as gallic acid equivalents (GAE) in milligrams per gram dry weight sample.

Determination of Total Antioxidant Activity

Antioxidant activity of methanolic extract of each part was assessed according to the method described by Brand *et al.* (1995) using the 1, 1-diphenyl-2-picrylhydrazyl (DPPH) radical-scavenging capacity with some modifications. Afterwards, 30 μL of methanolic extract of each part was mixed with 350 μL of DPPH (1mM in methanol) and methanol 100% was added to bring to the final volume of 2 mL. Following incubation of the mixture in the dark and room temperature for 20 min, the absorbance of the reaction mixture was read at 517 nm using a UV-Vis spectrophotometer. The volume of the sample extract required to cause a 50% decrease in the absorbance at 517 nm relative to the control (100%) was calculated. Methanol and ascorbic acid were used as the control and positive control, respectively. The inhibition ratio was calculated from the following equation:

Inhibition (%) = $\frac{[\text{Absorbance of the control} - \text{Absorbance of test sample}]}{\text{Absorbance of the control}} \times 100$

Extraction

The fresh tissues of treated plants were thoroughly ground with a cold mortar and pestle in an ice bath, until no fibrous residue could be seen. The grinding medium (500 mg) were completely extracted in 1,000 μL cool buffer phosphate (containing $\text{K}_2\text{HPO}_4 + \text{KH}_2\text{PO}_4$ 50 mM EDTA 0.5 mM and pH= 7) and centrifuged at 15,000 rpm at 5°C for 15 minutes. The supernatant was



used for determination of the superoxide dismutase (SOD) and peroxidase (POD) activity in the tissue.

Total Peroxidase

Peroxidase activity was assessed according to the method of Kalir *et al.*, (1984) with minor modifications. The reaction solution (1 mL) contained phosphate buffer (475 μ l, 50 mM; pH 7.0), guaiacol (475 μ L; 45 mM), H₂O₂ (475 μ L; 100 mM), and crude enzyme extract (50 μ L). Absorbance due to the formation of tetra-guaiacol was recorded at 470 nm using a UV-Vis spectrophotometer and enzyme activity was calculated as per its coefficient of extinction (26.6 mM⁻¹ cm⁻¹). One unit of enzymatic activity was defined as the amount of the enzyme that causes a change of 0.01 in absorbance per minute.

Total Superoxide Dismutase Activity

The activity of Superoxide dismutase was determined as the inhibition of the *p* - nitroblue tetrazolium chloride (NBT) photoreduction (Giannopolitis and Ries, 1977). The assay was conducted at 25°C in a total volume of one mL of 50 mM potassium phosphate buffer (pH 7.8), containing 0.1 mM EDTA, 1.3 μ M riboflavine, 13 mM methionine, 75 μ M NBT, and 50 μ L of enzymatic extract. Absorbance was monitored at 560 nm after 15 minutes of illumination, using a UV-Vis spectrophotometer. One unit of SOD is defined as the amount of enzyme that inhibits 50% of NBT photoreduction to blue formazan that monitored at 560 nm. SOD activity of the extracts was expressed as units of SOD per milligram of protein.

Analytical Methods and Statistical Analysis

The effect of three concentrations (1, 10 and 100 mg mL⁻¹) of each aqueous extract

was evaluated in the treated plants and growth criteria (germination, root length, shoot length, root fresh weight, shoot fresh weight, leaf number, shoot: root ratio) and biochemical parameters (POX and SOD) were assessed. The shoot:root ratio was calculated based on shoot and root fresh weight measurements. This experiment was carried out in factorial completely randomized design with four replicates and ten samples per replicate for all evaluated characters, except for POX and SOD which were done in one replicate. The treatment means were separated by Duncan's multiple range tests using SPSS 10.0.

RESULTS AND DISCUSSION

Germination and Growth Criteria

The results of Duncan's multiple range test showed that by increasing the concentration of root and shoot extract from 1 to 100 mg mL⁻¹ in the culture media, the growth criteria of treated plant including germination, root length, shoot length, root fresh weight, shoot fresh weight and leaf number were decreased, but the ratio of shoot to root was enhanced (Table 1). The most significant effect on germination and growth criteria of treated plants was obtained in 100 mg mL⁻¹ concentration that prevented germination and subsequent growth in most cases (Table 1). Goosefoot was highly resistant in the cases of fennel shoot, wormwood root and wormwood shoot extracts, even in the high concentration of 100 mg mL⁻¹ (Table 1). In other words, goosefoot was germinated and continued to grow in all extracts, except fennel root extract, in 100 mg mL⁻¹ concentration compared with lower concentrations (1 and 10 mg mL⁻¹) as the highest level of POX and SOD were observed in this concentration (Table 2).

At the same concentration (100 mg mL⁻¹), fennel and radish were not resistant to the goosefoot and redroot pigweed extracts, as in most treatments, germination and growth

Table 1. Duncan's multiple range tests comparing the mean of the effect of root and shoot extracts on growth criteria. ^a

Treated plant	Growth criteria	Root fresh weight (mg)			Shoot fresh weight (mg)			Shoot length (mm)			Leaf number			Shoot: root ratio			Number Germination					
		Concentration [mg.mL ⁻¹]									Extract											
		1	10	100	1	10	100	1	10	100	1	10	100	1	10	100	1	10	100			
Goosefoot	Fennel root	8.21 ^a	3.46 ^b	0 ^c	34.38 ^a	15.65 ^b	0 ^c	110.27 ^a	56.7 ^b	0 ^c	49.8 ^a	34.6 ^b	0 ^c	7.3 ^a	3.6 ^b	0 ^c	14.11 ^a	16.58 ^a	0 ^b	8.5 ^a	8 ^a	4 ^b
	Fennel shoot	9.28 ^a	7.05 ^b	4 ^c	35.52 ^a	27.8 ^b	15.8 ^c	115.2 ^a	94.78 ^b	60.02 ^c	51 ^a	45.4 ^b	4.15 ^c	7.65 ^a	7.1 ^{ab}	4 ^b	13.11 ^b	13.51 ^b	15.34 ^a	8.5 ^a	8.5 ^a	5.75 ^b
	Wormwood shoot	9.09 ^a	5.26 ^b	2.99 ^c	35.1 ^a	20.7 ^b	14.2 ^c	113.15 ^a	76.97 ^b	51.53 ^c	50.7 ^a	41.15 ^b	6.6 ^{ab}	7.35 ^a	6.6 ^{ab}	4 ^b	12.61 ^c	15.02 ^b	17.49 ^a	8.5 ^a	8.25 ^a	6 ^b
Fennel	Wormwood root	8.81 ^a	5.03 ^b	3.01 ^c	34.65 ^a	22.45 ^b	14.5 ^c	114.8 ^a	77.31 ^b	54.06 ^c	52.05 ^a	40.64 ^b	22.2 ^c	7.5 ^a	6.2 ^b	3.95 ^c	13.18 ^c	15.53 ^b	18.07 ^a	8.5 ^a	8.5 ^a	5.75 ^b
	Pigweed root	8.7 ^a	5.94 ^b	0 ^c	48.6 ^a	29.34 ^b	0 ^c	63.96 ^a	44.33 ^{ab}	0 ^b	74.21 ^a	49.63 ^b	0 ^c	3.86 ^a	2.98 ^b	0 ^c	7.56 ^a	7.51 ^a	0 ^b	9.33 ^a	8 ^b	0 ^c
	Pigweed shoot	8.71 ^a	6.06 ^b	0 ^c	49.45 ^a	31.46 ^b	0 ^c	64.65 ^a	44.03 ^{ab}	0 ^b	74.2 ^a	50.96 ^b	0 ^c	3.65 ^a	3.27 ^a	0 ^b	7.45 ^a	7.35 ^a	0 ^b	9.25 ^a	8 ^a	0 ^b
Radish	Goosefoot root	9.11 ^a	6.17 ^b	0 ^c	50.8 ^a	30.1 ^a	0 ^c	62.96 ^a	45.03 ^{ab}	0 ^b	72.27 ^a	51.33 ^b	0 ^c	3.53 ^a	3.2 ^{ab}	0 ^b	6.93 ^a	7.36 ^a	0 ^b	9.33 ^a	8 ^a	0 ^b
	Goosefoot shoot	9.27 ^a	7.52 ^b	3.1 ^c	50.13 ^a	40.8 ^b	9.85 ^c	65.36 ^a	49.8 ^b	25 ^c	73 ^{ab}	60.9 ^b	20.25 ^c	3.66 ^a	3.7 ^{ab}	2 ^b	7.06 ^b	6.66 ^b	8.32 ^a	8.66 ^a	8.25 ^a	6 ^b
	Pigweed shoot	15.71 ^a	10.24 ^b	0 ^c	27.2 ^a	18.07 ^b	0 ^c	184.27 ^a	126.39 ^b	0 ^c	45.25 ^a	38 ^b	0 ^c	3.75 ^a	3.10 ^b	0 ^c	12.51 ^a	12.17 ^a	0 ^b	5.5 ^a	3.75 ^b	0 ^c
Goosefoot	Pigweed root	14.72 ^a	6.72 ^b	0 ^c	28.06 ^a	11 ^b	0 ^c	185.91 ^a	103.98 ^b	0 ^c	45.51 ^a	28.12 ^b	0 ^c	2.96 ^a	2.6 ^{ab}	0 ^b	12.62 ^b	16.23 ^a	0 ^c	5 ^a	3.5 ^b	0 ^c
	Goosefoot root	15.35 ^a	10.38 ^b	0 ^c	27.75 ^a	19.28 ^b	0 ^c	183.91 ^a	127.5 ^b	0 ^c	48.05 ^a	38.41 ^b	0 ^c	3.39 ^a	3 ^{ab}	0 ^b	12.45 ^a	12.3 ^a	0 ^b	5 ^a	3.5 ^b	0 ^c
	Pigweed shoot	14.98 ^a	6.42 ^b	0 ^c	28.15 ^a	10.72 ^b	0 ^c	185.01 ^a	104.15 ^b	0 ^c	46.65 ^a	27.87 ^b	0 ^c	3.6 ^a	2.65 ^b	0 ^c	11.79 ^b	17.98 ^a	0 ^c	5.75 ^a	3.25 ^b	0 ^c

^a Different letters (a-c) indicate significant differences (Duncan's test at P<0.05 level).


**Table 2.** Peroxidase and superoxide dismutase activity of plant parts treated with different extracts.

Extract	Concentration (mg mL ⁻¹)	Treated plant	SOD (μm min ⁻¹ cc ⁻¹)		POX (U gr ⁻¹ FW)	
			Root	Shoot	Root	Shoot
Control	0	Goosefoot	165.48	41.73	28.62	2.36
Wormwood root	1	Goosefoot	283.19	87.58	56.73	4.34
Wormwood root	10	Goosefoot	909.57	116.04	131.78	5.01
Wormwood root	100	Goosefoot	1543.42	146.70	209.45	5.86
Wormwood shoot	1	Goosefoot	261.07	93.81	69.99	3.61
Wormwood shoot	10	Goosefoot	912.12	111.16	118.60	3.86
Wormwood shoot	100	Goosefoot	1484.62	159.37	187.52	4.93
Fennel root	1	Goosefoot	376.61	127.77	89.25	3.19
Fennel root	10	Goosefoot	1472.13	257.24	118.78	4.32
Fennel shoot	1	Goosefoot	301.08	68.34	27.43	2.97
Fennel shoot	10	Goosefoot	638.74	94.84	41.80	3.14
Fennel shoot	100	Goosefoot	763.59	189.99	72.02	4.84
Control	0	Fennel	285.20	99.51	7.18	1.48
Goosefoot root	1	Fennel	321.00	161.52	14.70	0.95
Goosefoot root	10	Fennel	412.93	182.27	18.86	2.51
Goosefoot shoot	1	Fennel	316.71	125.15	8.22	1.04
Goosefoot shoot	10	Fennel	374.02	163.77	10.43	1.90
Goosefoot shoot	100	Fennel	752.16	506.76	39.90	3.45
Redroot pigweed root	1	Fennel	336.82	154.85	16.73	1.63
Redroot pigweed root	10	Fennel	419.35	244.15	25.38	1.43
Redroot pigweed shoot	1	Fennel	323.43	160.55	29.08	0.79
Redroot pigweed shoot	10	Fennel	394.35	217.24	34.04	2.31
Control	0	Radish	194.71	114.91	92.60	9.07
Goosefoot shoot	1	Radish	437.82	168.12	174.59	17.83
Goosefoot shoot	10	Radish	726.55	206.32	387.93	26.37
Goosefoot root	1	Radish	393.55	129.58	226.70	17.12
Goosefoot root	10	Radish	712.37	220.22	417.38	23.24
Redroot pigweed root	1	Radish	272.90	128.57	280.61	14.47
Redroot pigweed root	10	Radish	637.15	264.187	461.43	16.61
Redroot pigweed shoot	1	Radish	793.96	176.24	368.66	109.69
Redroot pigweed shoot	10	Radish	921.02	327.20	593.97	101.02


of fennel and radish were prevented. In the present study, for all growth criteria, no significant difference was detected between lower concentrations of 1 mg mL⁻¹ and the control condition, while 100 mg mL⁻¹ almost inhibited the germination and subsequent growth (Table 1). Therefore, we focused on 10 mg mL⁻¹ concentration in order to determine their responses to the allelopathic effects.

Undoubtedly, germination is one of the most important plant growing stages and is severely affected by allelochemical components (Bogatek et al., 2006). No significant difference was observed between the effect of plants extracts and control conditions on germination at 10 mg mL⁻¹


concentration (Figure 1a-c), but at 100 mg mL⁻¹ concentration, root and shoot extracts severely decreased the seed germination (Table 1). In agreement with our results, many studies have reported the inhibitory effect of allelochemicals on germination. The effect of plant extract of *Nepeta meyeri* on germination and growth of seven plant species showed that with increasing the extract concentration, germination was reduced (Mutlu et al., 2010). Moreover, the allelopathic effects of two cultivars of sunflower (*Helianthus annuus* L.) at 2.5, 5, and 10% (m/v) concentration on germination of mustard (*Sinapis alba*) revealed that with increasing the extract concentration, seed germination was inhibited as decrease in



(a)



(b)



(c)

Figure 1. Comparison of the mean effect of 10 mg mL⁻¹ concentration of root and shoot extracts of (a) fennel and wormwood on goosefoot growth criteria, (b) redroot pigweed and goosefoot on fennel growth criteria and (c) redroot pigweed and goosefoot on radish growth criteria.. Different letters (a-d) indicate significant differences (Duncan's test at P< 0.05 level).



germination was significantly correlated with increased membrane deterioration (Bogatek *et al.*, 2006). Aqueous extract of plant parts at low concentrations had no effect on seed germination, but by increasing the concentration of the extract, seed germination displayed more reduction (Valera-Burgos *et al.*, 2012; Lin *et al.*, 2004).

Growth criteria of fennel, goosefoot, and radish at 10 mg mL⁻¹ concentration revealed that extracts severely reduced shoot length, fresh weight, root length and fresh weight, while shoot: root ratio was greatly increased compared to the control conditions (Figure 1a-c). Results showed that the highest effect on growth criteria were related to the fennel root extract (in the case of goosefoot), goosefoot root extract (in the case of fennel), and redroot pigweed shoot extract (in the case of radish). The highest amount of POX and SOD in both root and shoot of fennel, goosefoot, and radish at 10 mg mL⁻¹ concentration, were related to goosefoot root, fennel root, and redroot pigweed shoot extract, respectively. The least effect on growth criteria was related to the fennel shoot extract (in the case of goosefoot) and goosefoot shoot extract (in the case of fennel) (Table 2). The least amount of POX and SOD in both root and shoot of fennel and goosefoot at 10 mg mL⁻¹ concentration, were related to goosefoot and fennel shoot extract, respectively. Therefore, the goosefoot (in the case of fennel shoot extract) and fennel (in the case of goosefoot shoot extract) germinated and continued to growth at 100 mg mL⁻¹ concentration (Table 3). The effects of root and shoot extracts of wormwood (in the case of goosefoot) and goosefoot (in the case of radish) on growth were similar and no significant was observed between them.

In view of biochemistry, our results showed that root system was influenced more than shoot system (Table 2), as in all treated concentrations the increase in activities of POX and SOD of root were

more than that of shoot. Many studies have reported the inhibitory effect of allelochemicals on growth (Soltys *et al.*, 2011; Chowhan *et al.*, 2013; Mishra and Nautiyal, 2011). The effect of humic substance and its fractions were studied on morphology of *Arabidopsis thaliana* and results showed that humic fractions, especially at high concentrations, reduced root and shoot fresh weight and enhanced shoot: root ratio (Muscolo *et al.*, 2010). It is reported that shoot: root ratio of *Quercus rubra* was enhanced in the presence of *Dennstaedtia punctilobula* (Lyon and Sharpe, 1996). The effect of leaf extract of *Acacia pennatula* on germinability of three plant species revealed that general growth and especially the development of root in seedlings was reduced, reflecting shifting their biomass allocation model to a reduced root: shoot ratio which interferes in the development of the root system (Peguero *et al.*, 2012). Impairment of various metabolic activities due to plant leachate resulted in decreased root and shoot length (Singh *et al.*, 2009). Extract of *Pinus pinea* decreased root length of *Cistus* (Valera-Burgos *et al.*, 2012), which may influence seedling establishment in the field due to the differential access to water sources, hence, poor development of root could affect the ability of plant to cope with water deficiency condition, especially during summer.

Biochemical Parameters

The control treatments showed the maximum growth and minimum POX and SOD activity in both root and shoot compared to the other concentrations (Table 2). It was reported that the controls had the least POX activity and when plants were treated with different extract concentrations, POX and SOD were increased (Singh *et al.*, 2009; Bai *et al.*, 2009). Similarly, another study showed that by increasing the concentration of plant extract (*Eupatorium adenophorum*), POX and SOD activity of goosefoot and redroot pigweed leaves were

enhanced (Jinhu *et al.*, 2012). Allelochemicals cause oxidative damage, as evidenced by enhanced activity of ROS-scavenging enzymes and increased membrane lipid peroxidation (Lara-Nunez *et al.*, 2006). Moreover, antioxidative enzymes are the most important components in the scavenging system of ROS; and SOD is a major scavenger of superoxide, and its enzymatic action results in the formation of H_2O_2 . The H_2O_2 produced is then scavenged by catalase (CAT) and a variety of POX (Noctor and Foyer, 1998). Therefore, according to our results, POX and SOD systems of root and shoot of the plants encountered allelochemicals of different concentrations (even low concentration) and these systems were activated as a resistance mechanism in stressful condition in tissue culture media. Plants that germinated and continued to grow at high concentration of 100 mg mL^{-1} also had the highest POX and SOD activities (Table 2) Indicating further resistance.

According to the results, the plant parts containing more phenol have more antioxidant activity as wormwood shoot has the highest phenolic content and antioxidant activity (Total phenol= $57 \mu\text{g mg}^{-1}$ DW and DPPH IC50= 0.4 mg) (Table 3). In accordance to our results, previous researches (Caia *et al.*, 2004; Wojdylo *et al.*, 2007; Lu *et al.*, 2011) showed that there was a positive correlation between antioxidant activity and total phenolic content. Our data shows that different plant parts have different phenolic and antioxidative activity

and subsequently different allelopathic effects. For example, the root of fennel (in the case of goosefoot) and goosefoot (in the case of fennel) showed low phenolic content and, consequently, low antioxidant activity, revealing more allelopathic effects on growth criteria (Figure 1a and b) compared to fennel and goosefoot shoot with high amount of phenolic content and high antioxidant activity (Table 3). In addition, redroot pigweed shoot extract (in the case of radish) revealed low phenolic content and, consequently, low antioxidant activity (Table 3) and, therefore, more allelopathic effect on growth criteria than the other extracts (Figure 1c). More POX and SOD activity of parts (Table 2) accompanied by reducing growth criteria (Table 1). In accordance to our results, many researches have indicated that different plant parts contained different allelopathic effects (Sodaeizadeh *et al.*, 2009; Oueslati, 2003; Fernandez *et al.*, 2009). Some previous researchers have reported that phenolic components are the main cause of allelopathic effects (Garcia-Sanchez *et al.*, 2012; Li *et al.*, 2010; Jarchow and Cook, 2009), as more total phenolic content have more allelopathic effect on germination and subsequent growth (Ben-Hammouda *et al.*, 1995; Balezentiene and Seziene, 2010; Chon and Nelson, 2010). However, it is not the fact that all phenolic components have allelopathic effects (Inderjit 1996) and there is no simple relationship between total phenolic content and allelopathic effect (Yang *et al.*, 2009). These statements are in

Table 3. Total phenolic content and antioxidant activity of root and shoot.

Plant material	DPPH	Total phenol [$\mu\text{g.mg}^{-1}$ DW]
Wormwood shoot	IC50= 0.4 mg	57
Wormwood root	IC50= 0.53 mg	50
Fennel root	IC50= 12.53 mg	ND
Fennel shoot	IC50= 0.73 mg	38
Redroot pigweed shoot	IC50= 11.87 mg	ND ^a
Redroot pigweed root	IC50= 23.04 mg	ND
Goosefoot shoot	IC50= 0.8 mg	27
Goosefoot root	IC50= 23.64 mg	ND

^a No Detection.



agreements with our finding that plants containing more phenolic and antioxidative activity reveal less allelopathic effects.

CONCLUSIONS

Our finding explains that different plant parts have different allelopathic effects on plant species. In addition, more allelopathic effects due to increasing extract concentration lead to reducing growth criteria and increasing biochemical (POX and SOD) activity of each plant part. Germination inhibition depends on the extract concentration and confirms the importance of allelopathic effects of extract (especially high concentrations) as a mechanism for controlling seed germination. Therefore, tissue culture media could be an accurate and suitable tool for: (1) screening resistant plant and alien (non-endemic) plants that are not yet cultivated and no data are available about their resistance to prevalent weeds of the region, and (2) identifying high allelopathic plants as potential bioherbicide and invasive plant controller.

ACKNOWLEDGEMENTS

We acknowledge Iranian Biological Resource Center (IBRC) for funding this research.

REFERENCES

1. Anaya, A. L., Saucedo-García, A., Contreras-Ramos, S. M. and Cruz-Ortega, R. 2013. Plant-Mycorrhizae and Endophytic Fungi Interactions: Broad Spectrum of Allelopathy Studies. In: "Allelopathy: Current Trends and Future Applications", (Eds.): Cheema, Z. A., Farooq, M. and Wahid, A.. Springer-Verlag Berlin, Heidelberg, New York, Dordrecht, London, PP. 55-80.
2. Aryakia, E. and Hamidoghli, Y. 2010. Comparison of Kinetin and 6-benzyl Amino Purine Effect on in vitro Microtuberization of Two Cultivars of Potato (*Solanum tuberosum* L.). *Amer. Eur. J. Agri. Environ. Sci.*, **8**: 710-714.
3. Bai, R., Ma, F., Liang, D. and Zhao, X. 2009. Phthalic Acid Induces Oxidative Stress and Alters the Activity of Some Antioxidant Enzymes in Roots of *Malus prunifolia*. *J. Chem. Ecol.*, **35**: 488-494.
4. Balezentiene, L. and Seziene, V. 2010. Biochemical Impact of Dominant Extracts of Scots Pine Cuttings on Germination. *Polish J. Environ. Stud.*, **19**: 35-42.
5. Barazani, O. and Friedman, J. 2001. Allelopathic Bacteria and Their Impact on Higher Plants. *Crit. Rev. Microbiol.*, **27**: 41-55.
6. Ben-hammouda, M., Kremer, R. J., Minor, H. C. and Sarwer, M. 1995. Chemical Basis for Differential Allelopathic Potential of Sorghum Hybrids on Wheat. *J. Chem. Ecol.*, **21**: 775-786.
7. Bogatek, R., Gniazdowska, A., Zakrzewska, W., Oracz, K. and Gawronski, S. W. 2006. Allelopathic Effects of Sunflower Extracts on Mustard Seed Germination and Seedling Growth. *Biol. Plant.*, **50**: 156-158.
8. Bouman, H. and Tiekstra, A. 2001. Mineral Nutrition in Tissue Culture: Influence on Propagation and Quality of the Plantlets. Find Out How to Access Preview-only Content Plant Nutrition. *Developments Plant Soil Sci.*, **92**: 316-317.
9. Brand, W. W., Cuvelier, H. E. and Berset, C. 1995. Use of a Free Radical Method to Evaluate Antioxidant Activity. *Food Sci. Technol.*, **82**: 25-30.
10. Caia, Y., Luob, Q., Sunc, M. and Corke, H. 2004. Antioxidant Activity and Phenolic Compounds of 112 Traditional Chinese Medicinal Plants Associated with Anticancer. *Life Sci.*, **74**: 2157-2184.
11. Cheel, J., Tumova, L., Areche, C., Van Antwerpen, P., Neve, J., Zouaoui-Boudjeltia, K., San Martin A., Vokral, I., Wsol, V. and Neugebauerova, J. 2012. Variations in the Chemical Profile and Biological Activities of Licorice (*Glycyrrhiza glabra* L.), as Influenced by Harvest Times. *Acta Physiol. Plant.*, **35**: 1337-1349.
12. Chon, S. U. and Nelson, C. J. 2010. Allelopathy in Compositae Plants: A Review. *Agron. Sustain. Dev.*, **30**: 349-358.
13. Chowhan, N., Pal Singh, H., Batish, D. R., Kaur, S., Ahuja, N. and Kohli, R. K. 2013.

- β -pinene Inhibited Germination and Early Growth Involves Membrane Peroxidation. *Protoplasma.*, **250**: 691-700.
14. Cipollini, D., Rigsby, C. M. and Barto, E. K. 2012. Microbes as Targets and Mediators of Allelopathy in Plants. *J. Chem. Ecol.*, **38**: 714–727.
 15. Einhellig, F. A. 1999. Allelopathy: Current Status and Future Goals. *ACS Symp. Series*, **582**: 1-24.
 16. Einhellig, F. A. 1996. Interactions Involving Allelopathy in Cropping Systems. *Agron. J.*, **88**: 886-893.
 17. Elliott, L. F. and Cheng, H. H. 1987. Assessment of Allelopathy among Microbes and Plants. In: “*Allelochemicals: Role in Agriculture and Forestry*”. (Eds.): Waller, G. R.. American Chemical Society, Washington, DC, PP. 504–515.
 18. Fernandez, C., Monnier, Y., Ormeno, E., Baldy, V., Greff, S., Pasqualini, M. J. and Bousquet-Melou, A. 2009. Variations in Allelochemical Composition of Leachates of Different Organs and Maturity Stages of *Pinus halepensis*. *J. Chem. Ecol.*, **35**: 970–979.
 19. Garcia-Sanchez, M., Garrido, I., Casimiro, I. D. J., Casero, P. J., Espinosa, F., Garcia-Romera, I. and Aranda, E. 2012. Defence Response of Tomato Seedlings to Oxidative Stress Induced by Phenolic Compounds from Dry Olive Mill Residue. *Chemosphere*, **89**: 708–716.
 20. Giannopolitis, C. N. and Ries, S. K. 1977. Superoxide Dismutases. I. Occurrence in Higher Plants. *Plant Physiol.*, **59**: 309-314.
 21. Goodal, J., Witkowski, E. T. F., McConnachie, A. J. and Keen, C. 2012. Altered Growth, Population Structure and Realised Niche of the Weed *Campuloclinium macrocephalum* (Asteraceae) After Exposure to the Naturalised Rust *Puccinia eupatorii* (Pucciniaceae). *Biol. Invasions.*, **14**: 1947-1962.
 22. Inderjit. 1996. Plant Phenolics in Allelopathy. *Bot. Rev.*, **62**: 186-202.
 23. Jarchow, M. E. and Cook, B. J. 2009. Allelopathy as a Mechanism for the Invasion of *Typha angustifolia*. *Plant Ecol.*, **204**: 113–124.
 24. Jinhu, M., Guofang, X., Wenxiu, Y., Leilei, M., Mei, G., Yuguo, W. and Yuanhuai, H. 2012. Inhibitory Effects of Leachate from *Eupatorium adenophorum* on Germination and Growth of *Amaranthus retroflexus* and *Goosefoot glaucum*. *Acta Ecol. Sinica.*, **32**: 50–56.
 25. Kalir, A., Omri, G. and Poljakoff-Mayber, A. 1984. Peroxidase and Catalase Activity in Leaves of *Halimione portulacoides* Exposed to Salinity. *Physiol. Plant.*, **62**: 238–244.
 26. Kato-Noguchi, H., Thi, H., Sasaki, H. and Suenaga, K. 2012. A Potent Allelopathic Substance in Cucumber Plants and Allelopathy of Cucumber. *Acta Physiol. Plant.*, **34**: 2045–2049.
 27. Lara-Nunez, A., Romero-Romero, T., Ventura, J. L., Blancas, V., Anaya, A. L. and Cruz-Ortega, R. 2006. Allelochemical Stress Causes Inhibition of Growth and Oxidative Damage in *Lycopersicon esculentum* Mill. *Plant Cell Environ.*, **29**: 2009–2016.
 28. Li, Z. H., Wang, Q., Ruan, X., Pan, C. D. and Jiang D. A. 2010. Phenolics and Plant Allelopathy. *Mol.*, **15**: 8933-8952.
 29. Lin, D., Tsuzuki, E., Dong, Y., Terao, H. and Xuan, T. D. 2004. Potential Biological Control of Weeds in Rice Fields by Allelopathy of Dwarf Lilyturf. *Plants BioControl.*, **49**: 187–196.
 30. Lu, M., Yuan, B., Zeng, M. and Chen, J. 2011. Antioxidant Capacity and Major Phenolic Compounds of Spices Commonly Consumed in China. *Food Res. Int.*, **44**: 530–536.
 31. Lyon, J. and Sharpe, W. E. 1996. Hay-scented Fern (*Dennstaedtia punctilobula* (Michx.) Moore) Interference with Growth of Northern Red Oak (*Quercus rubra* L.) Seedlings. *Tree Physiol.*, **16**: 923-932.
 32. Mallik, M. A. B. and Williams, R. D. 2008. Plant Growth Promoting Rhizobacteria and Mycorrhizal Fungi in Sustainable Agriculture and Forestry. In: “*Allelopathy in Sustainable Agriculture and Forestry*”. (Eds.): Zeng, R. S., Mallik, A. U. and Luo, S. M.. Springer Science+Business Media, LLC, New York, PP. 321-345.
 33. Mishra, S. and Nautiyal, C. S. 2011. Reducing the Allelopathic Effect of *Parthenium hysterophorus* L. on Wheat (*Triticum aestivum* L.) by *Pseudomonas putida*. *Plant Growth Regulator J.*, **66**: 155–165.
 34. Muscolo, A., Sidari, M., Logoteta, B. and Panuccio, M. R. 2010. Carboxyl and



- Phenolic Humic Fractions Alter the Root Morphology in *Arabidopsis thaliana* Seedlings. *Fresenius Environ. Bull.*, 19: 12b.
35. Murashige, T. and Skoog, F. 1962. A Revised Medium for Rapid Growth and Bioassays with Tobacco Tissue Cultures. *Plant Physiol.*, **15**: 473–497.
 36. Mutlu, S., Atici, O., Esim, N. and Mete, E. 2010. Essential Oils of Catmint (*Nepeta meyeri* Benth.) Induce Oxidative Stress in Early Seedlings of Various Weed Species. *Acta Physiol. Plant.*, **33**: 943–951.
 37. Noctor, G. and Foyer, C. H. 1998. Ascorbate and Glutathione: Keeping Active Oxygen under Control. *Annu. Rev. Plant Physiol.*, **49**: 249–279.
 38. Oueslati, O. 2003. Allelopathy in Two Durum Wheat (*Triticum durum* L.) Varieties Agriculture. *Ecosys. Environ.*, **96**: 161–163.
 39. Peguero, G., Lanuza, O. R. and Save, R. 2012. Allelopathic Potential of the Neotropical Dry-forest Tree *Acacia pennatula* Benth.: Inhibition of Seedling Establishment Exceeds Facilitation under Tree Canopies. *Plant Ecol.*, **213**: 1945–1953.
 40. Singh, A., Singh, D. and Singh, N. B. 2009. Allelochemical Stress Produced by Aqueous Leachate of *Nicotiana plumbaginifolia* Viv. *Plant Growth Regulator J.*, **58**: 163–171.
 41. Singleton, V. L. and Orthofer, R. 1999. Analysis of Total Phenols and Other Oxidation Substrates and Antioxidants by Means of Folin-Ciocalteu Reagent. *Methods Enzymol.*, **299**: 152–178.
 42. Siril, E. A. and Joseph, N. 2013. Micropropagation of Annatto (*Bixa orellana* L.) from Mature Tree and Assessment of Genetic Fidelity of Micropropagated Plants with RAPD Markers. *Physiol. Mol. Biol. Plant.*, **19**: 147–155.
 43. Sodaiezadeh, H., Rafieiolhossaini, M., Havlik, J. and Damme, P. V. 2009. Allelopathic Activity of Different Plant Parts of *Peganum harmala* L. and Identification of Their Growth Inhibitors Substances. *Plant Growth Regulator J.*, **59**: 227–236.
 44. Soltys, D., Rudzinska-Langwald, A., Kurek, W., Gniazdowska, A., Sliwinska, E. and Bogatek, R. 2011. Cyanamide Mode of Action during Inhibition of Onion (*Allium cepa* L.) Root Growth Involves Disturbances in Cell Division and Cytoskeleton Formation. *Planta.*, **234**: 609–621.
 45. Sturz, A. V. and Christie, B. R. 2003. Beneficial Microbial allelopathies in the Root Zone: the Management of Soil Quality and Plant Disease with Rhizobacteria. *Soil Tillage Res.*, **72**: 107–123.
 46. Teerarak, M., Charoenying, P. and Laosinwattana, C. 2012. Physiological and Cellular Mechanisms of Natural Herbicide Resource from *Aglaia odorata* Lour. on Bioassay Plants. *Acta Physiol. Plant.*, **34**: 1277–1285.
 47. Tigrea, R. C., Silvab, N. H., Santosc, M. G., Hondad, N. K., Falcãoe, E. P. S. and Pereiraf, E. C. 2012. Allelopathic and Bioherbicidal Potential of *Cladonia verticillaris* on the Germination and Growth of *Lactuca sativa*. *Ecotoxicology and Environ. Safety*, **84**: 125–132.
 48. Tognetti, P. M. and Chaneton, E. J. 2012. Invasive Exotic Grasses and Seed Arrival Limit Native Species Establishment in an Old-Field Grassland Succession. *Biol. Invasion.*, **14**: 2531–2544.
 49. Valera-Burgos, J., Diaz-Barradas, M. C. and Zunzunegui, M. 2012. Effects of *Pinus pinea* Litter on Seed Germination and Seedling Performance of Three Mediterranean Shrub Species. *Plant Growth Regulator J.*, **66**: 285–292.
 50. Wojdylo, A., Oszmianski, J. and Czemerzys, R. 2007. Antioxidant Activity and Phenolic Compounds in 32 Selected Herbs. *Food Chem.*, **105**: 940–949.
 51. Yang, W. D., Liu, J. S., Li, H. Y., Zhang, X. L. and Qi, Y. Z. 2009. Inhibition of the Growth of *Alexandrium tamarense* by Algicidal Substances in Chinese Fir (*Cunninghamia lanceolata*). *Bull. Environ. Contam. Toxicol.*, **83**: 537–541.
 52. Xuan, T. D., Toyama, T., Dang Khanh, T., Tawata, S. and Nakagoshi, N. 2012. Allelopathic Interference of Sweet Potato with Cogongrass and Relevant Species. *Plant Ecol. J.*, **213**: 1955–1961.

ارزیابی اثرات آلوپاتیک برخی گونه های گیاهی در محیط کشت بافت به عنوان یک روش دقیق انتخاب گیاه مقاوم و غربال گری علف کشهای زیستی

ا. آریاکیا، م. ر. نقوی، ز. فرهمند، و س. ا. ح. شاهزاده فاضلی

چکیده

تکنیک کشت بافت گیاهی می تواند شرایط استریل و قابل کنترلی را فراهم کند که در آن اثر دقیق و مستقیم ترکیبات مختلف را بر شاخص های رشد و نمو گیاهان مورد بررسی قرار داد. در این پژوهش اثر عصاره آبی (۱، ۱۰، و ۱۰۰ mg.mL-1) ریشه و شاخه گیاهان سلمه (*Chenopodium album*) و (L.) تاج خروس (*Amaranthus retroflexus* L.)، رازیانه (*Foeniculum vulgare*) و درمنه (*Artemisia absinthium* L.) بر جوانه زنی و پارامترهای بیوفیزیکی گیاه در بستر کشت بافت مورد بررسی قرار گرفت. نتایج نشان داد عصاره ریشه رازیانه که تقریباً فاقد ترکیبات فنولی و متعاقباً توان آنتی اکسیدانی کم بود، بیشترین و شدیدترین تاثیر آلوپاتیک را به ویژه در غلظت ۱۰۰ mg.mL-1 روی علف هرز سلمه دارد که ممکن است ناشی از ترکیباتی با پتانسیل کنترل بیولوژیک علف هرز و مهاجم سلمه باشند. گیاه سلمه نسبت به عصاره شاخه رازیانه، ریشه و شاخه آرتمیسیا مقاوم است ولی رازیانه و تربچه، در غلظت زیاد و مشابه ۱۰۰ mg.mL-1، نسبت به عصاره ریشه و شاخه تاج خروس و سلمه مقاوم نیستند. در پاسخ به ترکیبات آلوپاتیک نسبت شاخه: ریشه افزایش یافت و بیشترین میزان پراکسیداز و سوپراکسید دیسموتاز نیز در ریشه وجود داشت. بین پتانسیل آلوپاتیک گیاهان و محتوای فنول و توان آنتی اکسیدانتی رابطه مستقیم وجود نداشت. نتایج نشان داد تاثیرات آلوپاتیک عصاره روی شاخص های بیوفیزیکی و بیوشیمیایی به سطح غلظت عصاره و نوع اندام گیاهی بستگی دارد. بنابراین بستر کشت بافت ابزار دقیق و مناسبی برای غربال گری گیاه مقاوم به ترکیبات آلوپاتیک علف های هرز، شناسایی گیاهان با پتانسیل آلوپاتیک زیاد؛ احتمالاً به عنوان علف کش بیولوژیک و کنترل کننده گیاهان مهاجم می باشد.